| Children <u>11 Years and Older</u> Need These Immunizations to Attend School in Arizona 2008-2009 School Year | | | | | |
|--|--|--|---|--|----------------------|
| Immunizations | 1 st Dose | 2 nd Dose | 3 rd Dose | 4 th Dose | 5 th Dose |
| (DTaP/DTP) Diphtheria, Tetanus & Pertussis, (DT) Diphtheria & Tetanus, (Td) Tetanus & Diphtheria | Required | Required | Required | Required if dose #1 was given before 12 months of age. | · |
| An interval of at least 4 weeks is required between the first doses of DTaP/DT/Td. An interval of 6 months is required between the last two doses. | | | | | |
| (Tdap) Tetanus, Diphtheria & Pertussis (New Requirement) | Required for 6 th grade students who are 11 years and older and have had 5 years pass since their last DTaP/DTP//Td dose. | | | | |
| Students 11 years and older in all grades are required to receive one dose of Tdap if 10 years have passed since their last dose of DTaP/DTP/DT/Td, or if they are in the process of receiving their first 3 doses of tetanus/diphtheria vaccine. Students who have received 1 dose of Tdap will need a Td booster after 10 years. | | | | | |
| (MV) Meningococcal Vaccine (New Requirement) | Required for 6 th grade students who are 11 years and older. | | | | |
| A dose of MV is recommended at 11 years of age. Because MV is licensed for use in children 2 years and older, doses given between 2 and 11 years of age are valid and acceptable. | | | | | |
| (IPV/OPV) Polio | Required | Required | Required | Not required if dose #3 was given at 4+years. 4 doses meet the requirement even if all were given in the first year of life. | |
| An interval of at least 4 weeks is required between all Polio doses. | | | | | |
| (MMR) Measles, Mumps, & Rubella | Required | Required | | | |
| MMR #1 is due at 12-15 months of age. MMR #2 may be given 1 month after MMR #1. MMR #2 is required for grades K-12. | | | | | |
| (Hep B) Hepatitis B | Required | Required | Required. Minimum age for final Hep B dose is 24 weeks. | Required only if dose #3 was given before 24 weeks of age. | |
| Hep B #1 may be given at birth. Dose #2 is due 4-8 weeks after dose #1. Dose #3 is due 2-5 months after dose #2, at least 4 months after dose #1 and no sooner than 24 weeks of age. A 2-dose adolescent series may be given to children 11-15 years of age over a 4-6 month period. | | | | | |
| <u>Varicella</u> (Chicken Pox) | Required | Required if dose #1 was given at 13+ years of age. | | | |
| In the 2008-2009 school year, children enrolled in preschool, kindergarten and grades 1, 2, 3, 4, 7, 8, 9 and 10 are required to be immunized against varicella (chicken pox) or have a history of chicken pox disease. Parental recall of chicken pox is acceptable. One dose of varicella vaccine meets the requirement for all children except those who receive varicella #1 at 13+ years of age. Children who receive the 1 st dose of varicella vaccine at 13+ years of age need 2 doses, spaced at least 1 month apart. Note: If MMR and Varicella vaccine doses were not given on the same day, they must be given at least 28 days apart. | | | | | |

Arizona Department of Health Services, Immunization Program Office, (602) 364-3632, Toll-free Number: (866) 222-2329.